

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Evaluation of Neural Regulation and Microglial Responses to Brain Injury in Larval Zebrafish Exposed to Perfluorooctane Sulfonate

Shannon E. Paquette, Nathan R. Martin, April Rodd, Katherine E. Manz, Eden Allen, Manuel Camarillo, Hannah I. Weller, Kurt Pennell, and Jessica S. Plavicki

Table of Contents

Table S1. Survival rate of zebrafish larvae exposed to PFOS over time.

Table S2. Primers used for qRT-PCR and genotyping.

Table S3. PFOS body burden in zebrafish larvae.

Table S4. Significantly enriched metabolic pathways following 28 μ M PFOS exposure in 3 dpf larvae.

Figure S1. Microglia quantification, cell death, and proinflammatory cytokine mRNA analysis in PFOS-exposed larvae.

Figure S2. Control for the optogenetic microglia stimulation experiment by stimulating halorhodopsin negative microglia.

Figure S3. Functional neuroimaging to understand the effects of toxicant exposure on neuronal activity.

Figure S4. Regional and global brain morphology in larvae exposed to varying concentrations of PFOS.

Figure S5. MWAS of heads collected from control versus PFOS-exposed larvae.

Figure S6. Light/dark behavioral assay in larvae exposed to varying concentrations of PFOS.

Figure S7. Absolute and relative time spent in center throughout the light/dark behavioral response assay.

Figure S8. CaMPARI analysis of the habenula in larvae exposed to varying concentrations to PFOS.

Figure S9. Distance traveled in control versus PFOS-treated wildtype and macrophage mutant larvae.

Figure S10. Microglia response to brain injury in control versus PFOS-exposed versus PTZ-exposed larvae at 3 dpf.

Figure S11. Light/dark behavioral analysis in PFOA-exposed larvae.

Figure S12. Body length of 5 dpf exposed to PFOA.

References

Additional File- Excel and Video Document